

An Indication From the Magnitude of CP Violations That Gravitation Is A Possible Cause of Wave-Function Collapse

Daniel I. Fivel

*Department of Physics, University of Maryland
College Park, MD 20742*

(October 8, 1997)

We consider experimental evidence for the hypothesis that the Planck energy, $E_p \approx 10^{19} \text{ GeV}$, sets the scale ϵ at which wave function collapse causes deviations from linear Schrödinger evolution. With a few plausible assumptions about the collapse process, we first show that the observed CP violation in K_L decay implies a lower bound on ϵ remarkably close to E_p . If the bound is saturated, the entire CP violation is due to collapse and a prediction made that the branching ratio for CP violation in the B meson decay will be $\gamma \approx 10^{-5}$. We then show that the assumptions are consequences of a simple non-linear, stochastic modification of the Schrödinger equation with ϵ setting the scale of the non-linearity.

In his *Lectures on Gravitation* [1], R.P. Feynman suggested that we at least consider the possibility that gravity need not be quantized because it is actually responsible for a deviation from linear Schrödinger evolution at the Planck scale, $E_p \approx 10^{19} \text{ GeV}$. This is not implausible since a particle with mass larger than $M_p = E_p/c^2 \approx 10^{-5} \text{ gm}$ has a Compton wavelength smaller than its Schwarzschild radius.

Feynman also observed that because the Planck mass M_p is essentially macroscopic, such a deviation might explain the apparent instability and collapse of Schrödinger cat states in which there is dispersion of a macroscopic observable. Indeed, the measurement problem of quantum mechanics arises from the absence of a generalization of the Schrödinger equation that interpolates between linear evolution and collapse. Such a generalization requires a universal energy scale ϵ to which a dispersion Δ arising from entanglement can be compared in determining whether the collapse time τ_c is large or small in relation to the time scale $\hbar/\Delta$ of linear Schrödinger evolution. The question then is whether E_p supplies that scale. Since there is, as far as we know, no interaction weaker than gravity, the establishment of a *lower bound* for ϵ in the vicinity of E_p would lend strong support to the candidacy of gravity as the cause of collapse.

The simplest guess we can make from dimensional analysis is

$$\tau_c = \hbar \epsilon / \Delta^2. \quad (1)$$

If collapse does not actually happen, as some believe, we would have $\tau_c \rightarrow \infty$, i.e. $\epsilon \rightarrow \infty$. If collapse does happen, $\epsilon \approx E_p$ implies that collapse is very slow and difficult to measure.

To obtain a lower bound experimentally one might consider the possibility of detecting collapse by its effect on the observable EPR correlations of entangled pairs. But in today's technology such experiments are only conceivable in the realm of quantum optics where $\Delta \approx 1 \text{ eV}$. Absence of observed collapse would then imply $\epsilon \gg 1 \text{ eV}$. Even if such measurements could be made with hadrons, our bound will only move up to 1 GeV, and we are still nineteen orders of magnitude too low.

Fortunately, however, there are entanglement situations involving elementary particles in which a conspiracy of circumstances provides an opportunity to detect a much larger lower bound. Suppose we have an entangled state which is unstable, decays in time τ , and has an observed branching ratio $\gamma \ll 1$ for breaking a symmetry. If τ_c were not very much larger than τ , there would be time to first collapse into the factorized constituents of the entangled state. If the branching ratio for violation in such case is $\gamma_c \gg \gamma$, a lower bound on τ_c follows. If collapse obeys an exponential law one obtains:

$$\tau_c \geq \gamma_c \tau / \gamma \implies \left(\frac{\epsilon}{\Delta} \right) \geq \left(\frac{\tau \Delta}{\hbar} \right) \left(\frac{\gamma_c}{\gamma} \right), \quad (2)$$

with saturation indicating that all of the violation is coming from the collapse channel.

Thus we can obtain a lower bound on ϵ much larger than Δ by having either or both of the following: (I) $\tau \Delta / \hbar \gg 1$, i.e. an unusually long lifetime because of the violation of a good conservation law. (II) γ / γ_c small, i.e. a small observed symmetry breaking but one that would be large if collapse happens before decay.

Both of these ingredients are present in the CP violating decay of the K_L meson which in the quark model is the state:

$$K_L = \alpha|s, 1\rangle|\bar{d}, 2\rangle + \beta|d, 1\rangle|\bar{s}, 2\rangle, \quad \alpha = 1/\sqrt{2} = -\beta. \quad (3)$$

This is a superposition of the K_o and $\bar{K}_o$ which are the summands on the right, but it is an *entanglement* of the two-level di-quark system indicated by label 1 and the two-level di-antiquark indicated by label 2. As will be made clear below, the relevant value of the dispersion Δ driving collapse is the total of the dispersion in each of the two entangled systems *separately*. Thus it is the dispersion arising from the mass difference $\delta m \approx 200$ MeV of the s and d quarks that we must use for Δ not the mass difference of the K_o and $\bar{K}_o$ which is zero. It is this large value together with the very long life of the K_L , $\tau \approx 5 \times 10^{-8} s$, resulting from a combination of strangeness changing and small phase space that supplies the principal amplification factor (I). The amplification factor (II) results from having $\gamma_c = 0.5$ because the K_o and $\bar{K}_o$ are not CP eigenstates, together with a small measured CP branching ratio, $\gamma \approx 2 \times 10^{-3}$. Putting these numbers into (2) we obtain the remarkable result:

$$\epsilon \geq E_p/8\pi. \quad (4)$$

We cannot, of course, rule out the possibility that ϵ is much larger, or even infinite. If so this occurrence of a bound comparable to E_p and *twenty eight orders of magnitude higher than any present experiment could have anticipated* is just a bizarre coincidence. Absent such a coincidence we are forced to conclude that collapse does indeed “happen” and that gravitation is somehow implicated. It also suggests, but does *not* require, an economical explanation of CP violations, namely that the bound is saturated, thus making collapse the sole cause.. This would mean that the CKM phase is zero. Moreover in the B-system, insertion of the b-d mass difference of 5 GeV for Δ and the lifetime $10^{-12} s$ for τ then predicts a branching ratio of $\gamma \approx 10^{-5}$.

Assuming that (4) is indeed a manifestation of a dynamical collapse mechanism, we are motivated to carry out the task that will occupy the remainder of this paper, namely the construction of a generalization of Schrödinger dynamics with one universal energy parameter ϵ whereby one can *deduce* the various ingredients of the above argument i.e. the validity of (1), the exponential law for collapse, and the independent contribution to Δ from the energy dispersions in each constituent of the entangled state. We shall arrive at the generalization through a sequence of steps intended to show that where choices are available we have made the simplest consistent with physical constraints.

We know that the modified Schrödinger equation must be non-linear to allow collapse to happen. But we also know [2–4] that non-linearities must be stochastic (noisy) or else pathologies will appear, i.e. the non-linearity can be exploited to send super-luminal signals and reduce the entropy of closed systems.

To keep the notation uncluttered we develop the theory for two-level entangled states such as (3) We will assume that the two levels for particle-1 are $\pm E$ which can be done by adding a multiple of the unit matrix to the Hamiltonian matrix. We first develop the equation for the case where only one of the particles has significant energy dispersion and then extend it to allow contributions from both. We will thus take the Hamiltonian to be:

$$H_{tot} = H \otimes I, \quad H = E\sigma, \quad \sigma = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (5)$$

We obtain a first clue as to the kind of stochastic non-linearity we need by looking at wave-function collapse thermodynamically. It is a peculiar irreversible process in which an entangled pure state such as (3), is driven into another pure state which is factorized. As normally defined all pure states have zero entropy. But one notes that entangled states contain more information than do pure states. For if a measurement is made on particle-1, paying no attention to particle-2, the density matrix becomes the reduced density matrix whose entropy is larger, whereas for a factorized state there would be no change. Thus (3) has a negative entropy of entanglement

$$S(x) \equiv x \log x + (1 - x) \log(1 - x), \quad x = |\alpha|^2, \quad (6)$$

which rises to zero when collapse takes place. We thus expect that the non-linear term that we add to the Schrödinger equation to induce collapse should depend on the wave function in such a way that its strength is correlated with the magnitude of this entanglement entropy.

There is an elegant and natural way to introduce non-linearities into the Schrödinger equation. If Ψ is an n-component spin wave function, and H is an $n \times n$ matrix then

$$\hbar \frac{d\Psi}{dt} = -iH\Psi \quad \Longleftrightarrow \quad \hbar \frac{d\Psi}{dt} = -iH \frac{\partial}{\partial \Psi^*} |\Psi|^2, \quad |\Psi|^2 = \sum_j |\Psi_j|^2. \quad (7)$$

This is the so-called “geometric formulation of quantum mechanics” [5,6] which makes the Schrödinger equation exhibit the (Kähler) symplectic structure [7,8] of Hamilton’s equations for conjugate dynamical variables Ψ and Ψ^* . One may now look for suitable modifications of the Schrödinger equation by adding something to $|\Psi|^2$ which is more than quadratic in the Ψ ’s and is thus like adding anharmonic terms to a classical oscillator. Our problem is to find such a quantity that is correlated with the entanglement entropy and is expressed simply in terms of the Ψ ’s.

We obtain a second clue as to the kind of quantity we need from the observation that entanglement is invariant under one-particle unitary transformations, i.e.

$$|\Psi\rangle = \sum_{j,k} \psi_{jk} |j, 1\rangle |k, 2\rangle \rightarrow (U \otimes V) |\Psi\rangle. \quad (8)$$

We can construct invariants under such transformations by observing that there is a natural map from $|\Psi\rangle$ to the operator

$$\psi = \sum_{j,k} \psi_{jk} |j, 1\rangle \langle k, 2|, \quad (9)$$

which is mapped under $U \otimes V$ to $U\psi V^\dagger$. Thus functions of the traces of powers of $\psi^\dagger\psi$ are invariants as is the determinant

$$\mathcal{E}(\psi) \equiv \text{Det} (\psi^\dagger\psi). \quad (10)$$

For the situation in which we are interested, ψ is a two by two matrix so that all of the invariants are functions of $\text{Tr} (\psi^\dagger\psi)$ and $\mathcal{E}$. Since $\text{Tr} (\psi^\dagger\psi)$ is just the norm $|\Psi|^2$ in (7) which gives the linear Schrödinger equation, we are led to consider whether $\mathcal{E}$ might be the quantity we are seeking.

In the case of the state (3) we have $\mathcal{E} = |\alpha|^2|\beta|^2 = x(1-x)$ so that it is correlated with the entanglement entropy (6) in the right way, having its maximum when the $|S|$ is largest and falling to zero when $|S| = 0$. That it is a quite general measure of entanglement can be seen from the following argument: What characterizes entanglement is the inability to make a prediction about the behavior of one constituent independently of its partner. Thus in a completely entangled state a constituent can be found with equal likelihood in any state if no measurement is made on its partner. In contrast, for a system in the factorized state $|1\rangle|2\rangle$, it is certain that particle -1 will not be found in a state orthogonal to $|1\rangle$. Now one sees that a necessary and sufficient condition for there to be zero probability of finding a particle in some state when no measurement is made on the other, is that such a state be an eigenstate of $\psi^\dagger\psi$ with eigenvalue zero. There is an eigenstate with eigenvalue zero if and only if $\mathcal{E}$ vanishes.

It appears then that we should be able to build our modified dynamics by rewriting (7) in terms of ψ and replacing $|\Psi|^2$ by a suitable combination of $\text{Tr} (\psi^\dagger\psi)$ and $\mathcal{E}$. Taking advantage of the identity

$$\text{Det}(I + \nu A) \equiv I + \nu \text{Tr}(A) + \nu^2 \text{Det}(A), \quad (11)$$

valid for any 2×2 matrix A and scalar ν , we are led to consider an equation of the form

$$\frac{\hbar}{\epsilon} \frac{d\psi}{dt} = \pm \sigma \frac{\partial}{\partial \psi^*} \text{Det} (1 \mp i\eta \psi^\dagger\psi), \quad \eta = E/\epsilon, \quad (12)$$

in which we have one new universal parameter ϵ . Expanding the determinant by (11) and working out the partial derivatives one obtains:

$$\hbar \frac{d\psi}{dt} = -iH\psi \pm \eta \text{Det} (\psi^\dagger\psi) H\psi^{\dagger-1}, \quad (13)$$

which shows that the equation reduces to the ordinary Schrödinger equation plus a nonlinear term that disappears if we let $\epsilon \rightarrow \infty$. This would be the case if collapse did not really happen at all.

The linear Schrödinger equation has a time reversal symmetry that shows up in its invariance under $t \rightarrow -t$ and $\psi \rightarrow \psi^*$. The non linear term which has no i does not preserve this symmetry. With a random sign in (13) the symmetry can be preserved as an ensemble average. This suggests a simple and natural way to introduce the noise: *We assume that during the evolution of the state of a particle, the signs in (13) fluctuate randomly.*

The description of the noise is not complete, of course, until we also specify the intervals between the random sign fluctuations. Before doing so let us examine the solution of the equation for a *fixed* choice of sign. Because of the

invariance of the determinant, the linear term in (13) is transformed away by going to the interaction picture, i.e. by the transformation $\psi \rightarrow e^{-iHt}\psi$. The ψ -matrix representing the state (3) will be initially diagonal and, because of the absence of coupling, will remain diagonal. We thus arrive at a pair of coupled differential equations for $x = |\alpha|^2$ and $y = |\beta|^2$, namely:

$$\frac{dx}{dt} = -\frac{xy}{\tau_c}, \quad \frac{dy}{dt} = \frac{xy}{\tau_c} \quad (14)$$

with τ_c given by (1), and $\Delta = E\sqrt{2}$. One observes that $x + y$ is conserved, and the process *terminates* when either x or y vanishes. The solution for x is seen to be: $x = 1/(1 + e^{t/\tau_c})$ which is essentially exponential with characteristic time τ_c .

We now introduce the noise in the manner suggested above by letting the signs in (13) fluctuate randomly. Writing $dx = \pm\delta$, $dy = \mp\delta$ with $\delta = xydt/\tau_c$ one sees that if the fluctuations occur in such a way that δ is held fixed the process is a random walk with boundary, sometimes called the “gambler’s ruin” [10,11]. One thinks of x and y as the fortunes of two gamblers who bet δ on each toss of a coin, the victor being indicated by the sign. The game ends when one is wiped out. This game is known to be a *martingale* i.e. the probability of winning is proportional to one’s initial fortune. Hence the probability for $x \rightarrow 1$ is x and for $y \rightarrow 1$ is y which is just the quantum mechanical collapse rule *but now produced dynamically rather than as a separate postulate*.

However, this stochastic process is too noisy and will result in an infinitely long process when we let dt go to zero. For if the stake gets small the weaker player can have runs of luck so that the average length of the game increases without limit. The way around this is to let the “game” be played as the game of “double or nothing” [9] in which the weaker of the two players bets the full amount of his fortune on any play. This game is also a martingale [12], but its average length is two (or 1 if $x = y = 1/2$).

As noted earlier we only considered the effect of the energy dispersion arising from one of the two particles in the entangled state. Effects of the second particle are included by adding a second term on the right of (12) identical to the first but with σ acting from the right on the gradient and the E value appropriate to the second particle. In general there need be no correlation between the signs of the noise in the two terms, and the result will be to simply add the dispersions. The fact that it is the sum of the energy dispersions that is relevant rather than the dispersion of the total energy (which is zero) is easily seen physically by the example of an entangled state consisting of a live cat A and dead cat B with dead cat A and live cat B. There is no dispersion of the total temperature, but the state must clearly collapse rapidly due to the elevated free energy associated with the dispersion of the temperatures in each cat separately.

We have now produced a stochastic, non-linear modification of the Schrödinger equation that interpolates between linear deterministic evolution and non-linear, stochastic collapse and from which we could derive (1) and the exponential collapse law with no stochastic tails.. Our final task is to show that our modified Schrödinger equation enjoys the same “peaceful coexistence” with the other basic laws of physics that the linear Schrödinger equation enjoys. This is particularly important because we know [2–4] that the noise is essential for this.

(1) The second law of thermodynamics: Consider the change in the average entanglement of an ensemble with a given x resulting from a noise fluctuation. We have

$$0.5(x + \delta)(1 - x - \delta) + (x - \delta)(1 - x + \delta) - x(x - 1) = -\delta^2 \quad (15)$$

Thus the average entanglement always decreases which means that the entropy always increases. Since the maximum average entanglement decrease is achieved for the largest δ consistent with the given x which is the minimum of x and $1 - x$ we see that *the “double or nothing” noise rule is equivalent to the requirement that entropy increase at the maximum possible rate*.

(2) Conservation of energy: One may calculate $\frac{d}{dt} \text{Tr}(\psi^\dagger H \psi)$ from (13). Noting that the total time for the process is $\approx 2\tau_c$ one obtains an upper bound on the energy fluctuation. Comparing this with the dispersion, one finds that the fluctuation lies within the uncertainty principle limit.

(3) Causality: The random sign fluctuations reproduce the collapse postulate of conventional quantum mechanics, and so the outcome of measurements has the same unpredictability. But it is this unpredictability that prevents capitalizing on EPR correlations to send super-luminal messages. Thus the status of the non-linear, stochastic theory vis a vis causality is the same as that of the linear equation plus collapse postulate. There are grounds for optimism that one can develop an explicitly covariant version because of the invariance of $\text{Det}(\psi^\dagger \psi)$ under all possible one-particle unitary transformations.

Since we have proposed a resolution of the measurement paradox, let us point out its salient differences from other proposed resolutions [13–15]. As in other dynamical theories, collapse actually happens, rather than merely appearing to happen. In this respect it differs from relative state and modal interpretations. [16] The noise is internal so it also happens in closed systems, a feature missing in the so-called [17] FAPP (“for all practical purposes”) approaches which rely on external noise reservoirs to produce decoherence. Finally it differs from spontaneous localization theories by denying favoritism to any direction in Hilbert space, but rather relying on the entanglement and thermodynamics to decide the direction of collapse by lowering the free energy. Having noise fluctuations occur at intervals influenced by the stochastic process itself we avoid the necessity of having external “hits” [18] and the necessity of having more than one fundamental parameter. If ϵ is simply related to E_p as the bound (4) suggests, there will be *no* new parameters in our modified equation.

The theory presented above is restricted to the simplest dynamical situation, that of two systems, each with two levels. Since all measurements in principle can be reduced to yes-no questions, this treatment is adequate for demonstrating how the measurement paradox may be resolved. However, to make it into a complete theory we must extend the non-linear equation to systems of many particles and to infinite dimensional Hilbert spaces. We anticipate that the thermodynamic view of collapse will be extremely useful for this, because the optimal decreasing of entanglement free energy becomes a physical basis by which the system “decides” along which basis in Hilbert space to collapse.

When nearing the conclusion of this paper the work of L. Hughston [19] was brought to my attention. He has clearly recognized that (1) will be obtained in stochastic evolution, and his non-linear stochastic equation also has the symplectic geometric form. In considering the possibility that the Planck scale may be involved he notes the relative accessibility of the combination $\hbar E_p \approx 8MeV^2s$. His analysis of the geometric content of stochastic dynamics will no doubt be invaluable in seeking to generalize the theory to more complex systems. It might also provide machinery for attacking the puzzle of why gravitation is playing the role it seems to be playing in the dynamics of wave function collapse.

Acknowledgements I wish to thank A. Dragt, A. Jawahery, A. Shimony, G. Snow, and J. Sucher for useful discussions. I also owe a profound debt of gratitude to T. Jacobson for much insightful criticism at various stages of this work, and most importantly for pointing out the relationship of the entanglement measure to the relative entropy.

-
- ¹ R. P. Feynman, *Lectures on Gravitation*, p.12,13, Addison-Wesley, Reading 1995
 - ² N. Gisin, *Helv. Phys. Acta* **62** (1989) 363
 - ³ A. Peres, in *Complexity, Entropy, and the Physics of Information*, SFI Studies in the Science of Complexity, vol. VIII, Ed. WH. Zurek, Addison-Wesley, 1990, p. 345
 - ⁴ J. Polchinski, *Phys. Rev. Let.* **66**, 397 (1991)
 - ⁵ T. W. B. Kibble, *Commun. Math. Phys.* **65**, 189 (1979).
 - ⁶ A. Heslot, *Phys. Rev.* **D31**, 1341 (1985)
 - ⁷ L. P. Hughston, in *Twistor Theory*, Ed. S.A. Huggett, Marcel Dekker, (1995)
 - ⁸ A. Ashtekar and T. A. Schilling gr-quant-th 9706069
 - ⁹ D. I. Fivel, *Phys. Rev. A*, **56** (1997) 146
 - ¹⁰ P. Pearle, *Phys. Rev. A* **39** (1989) 227
 - ¹¹ W. Feller, *An Introduction to Probability Theory and Its Applications* p. 234 Vol. 1”, John Wiley & Sons
 - ¹² A. P. Maitra and W. D. Sudderth, *Discrete Gambling and Stochastic Games*, Springer 1996.
 - ¹³ D. Z. Albert, *Quantum Mechanics and Experience* (Cambridge University Press 1992)
 - ¹⁴ A. J. Leggett in “*Quantum Implications*”, Ed. B. J. Hiley and F. David Peat, Routledge, N.Y (1987) Ch. 5
 - ¹⁵ A. Shimony, in *Search for a Naturalistic World View, Vol. II*. Cambridge University Press (1993)
 - ¹⁶ J. Bub, *Interpreting the Quantum World*, Cambridge University Press, (1997)
 - ¹⁷ J. S. Bell, in *Sixty-Two Years of Uncertainty*, Ed. A. I. Miller, NATO ASI Series, Plenum Press, N.Y., (1989)
 - ¹⁸ G. C. Ghirardi, A. Rimini, and T. Weber, *Phys. Rev.* **D34** (1986) 470
 - ¹⁹ L. P. Hughston, *Proc. R. Soc. Lond.* **A452**, 953 (1996)